

## Testicular Fibrous Pseudotumor: The Role of Ultrasound Examination and Intraoperative Frozen Section to Prevent Unnecessary Orchiectomy


To the Editors:

Six years after the publication of the last case of inflammatory testicular pseudotumor,<sup>1</sup> we have a further case to add to the overview of the pseudo-neoplastic entities arising in the testis and paratesticular structure,<sup>2</sup> particularly interesting for its clinical management.

A 70-year-old man presented with an asymptomatic, palpable nodule at the right testis. Ultrasonography (US) showed an 8-mm, solid, vascularized lesion at the testicular surface (Figure 1A). Testicular markers were normal.

The lesion was therefore excised (Figure 1B): its benign nature was confirmed at frozen section and the testicle spared (Figure 1C). Final microscopic examination disclosed a well-outlined, fibrous-like nodule composed of hyalinized collagen bands (Figure 1D). Normal findings were seen in a 6-month US follow-up.

In the literature, fibrous pseudotumors of the tunica vaginalis were first described by Balloch in 1904.<sup>3</sup> They are rare, benign, reactive lesions of the testicular tunica that can affect every age, although the third decade has a peak incidence.<sup>4</sup> Physical examination, US, and serum markers are the standard evaluation.<sup>5</sup> Fibrous pseudotumors have been described as hypoechoic nodules arising from the testicular surface at US. Since fibrous pseudotumors



**Figure 1**

(A) Simultaneous US and palpation revealed that the lesion was surrounded by a small quantity of fluid and arose from the testicular surface, with a small stalk containing a feeding vessel (arrow). (B) The excised lesion with its pedunculated stalk (arrow). (C-D) Hematoxylin-eosin-stained specimen (C, 2 $\times$ ; D, 40 $\times$ ) illustrating a well-circumscribed nodule composed of hyalinized collagen bands with interspersed lymphocytes. Nuclear atypia and mitoses are absent.

may mimic malignancies, radical orchiectomy has often been performed.<sup>6</sup> It is our purpose to underline the importance of US examination with simultaneous palpation and intraoperative frozen section to prevent unnecessary orchiectomy.

**Alberto Olivero, M.D.**

**Michela Campora, M.D.**

**Carlo Toncini, M.D.**

**Aldo F. De Rose, M.D.**

**Lorenzo E. Derchi, M.D.**

**Carlo Terrone, M.D.**

**Valerio Vellone, M.D., Ph.D.**

**Bruno Spina, M.D.**

From the Department of Urology, University Hospital Policlinico San Martino-IRCC; the Department of Surgical and Diagnostic Sciences (DISC), University Unit of Pathology, University of Genova; the Department of Anatomical Pathology, Hospital Policlinico San Martino-IRCC; and the Department of Health Sciences (DISSAL), University of Genova, Genoa, Italy.

Drs. Olivero and Campora contributed equally.

Address correspondence to: Michela Campora, M.D., Department of Integrated Surgical and Diag-

nostic Sciences, University of Genova, Via de Toni 14, 16132 Genoa, Italy (mikycampora@yahoo.it).

### References

1. Torres Gomez FJ, Fernández Machín P, Garcia Suarez RM: Right orchiectomy: Inflammatory testicular pseudotumor. *Arch Esp Urol* 2012;65:641-642
2. Algaba F, Mikuz G, Boccon-Gibod L, Trias I, Arce Y, Montironi R, Egevad L, Scarpelli M, Lopez-Beltran: Pseudoneoplastic lesions of the testis and paratesticular structures. *Virchows Arch* 2007;451:987-997
3. Balloch EA: Fibromata of the tunica vaginalis. *Ann Surg* 1904;39:396-402
4. Germaine P, Simerman LP: Fibrous pseudotumor of the scrotum. *J Ultrasound Med* 2007;26:133-138
5. Zhang Z, Yang J, Li M, Cai W, Liu Q, Wang T, Guo X, Wang S, Liu J, Ye A: Paratesticular fibrous pseudotumor: A report of five cases and literature review. *Front Med* 2014;8:484-488
6. Başal Ş, Malkoç E, Aydur E, Yıldırım İ, Kibar Y, Kurt B, Göktaş S: Fibrous pseudotumors of the testis: The balance between sparing the testis and preoperative diagnostic difficulty. *Turk J Urol* 2014;40:125-129

*Financial Disclosure:* The authors have no connection to any companies or products mentioned in this letter.

**Keywords:** frozen section, orchiectomy, palpation, testes, testicular fibrous pseudotumor, ultrasonography, ultrasound. (*Anal Quant Cytopathol Histopathol* 2018;40:106-107)